

Technical report bds:99-08

On the ultimate behavior of the sequence of consecutive powers of a matrix in the max-plus algebra*

B. De Schutter

If you want to cite this report, please use the following reference instead:

B. De Schutter, "On the ultimate behavior of the sequence of consecutive powers of a matrix in the max-plus algebra," *Linear Algebra and Its Applications*, vol. 307, no. 1–3, pp. 103–117, Mar. 2000.

Control Systems Engineering
Faculty of Information Technology and Systems
Delft University of Technology
Delft, The Netherlands
phone: +31-15-278.51.19 (secretary)
fax: +31-15-278.66.79
Current URL: <http://www.dcsc.tudelft.nl>

*This report can also be downloaded via http://pub.deschutter.info/abs/99_08.html

On the ultimate behavior of the sequence of consecutive powers of a matrix in the max-plus algebra

Bart De Schutter*

Abstract

We study the sequence of consecutive powers of a matrix in the max-plus algebra, which has maximum and addition as its basic operations. If the matrix is irreducible then it is well known that the ultimate behavior of the sequence is cyclic. For reducible matrices the ultimate behavior is more complex, but it is also cyclic in nature. We will give a detailed characterization of the rates and periods of the ultimate behavior for a general matrix.

Keywords: max-plus algebra, sequence of matrix powers, ultimate behavior

1 Introduction

We consider the sequence of consecutive powers of a matrix in the max-plus algebra, which has maximum and addition as basic operations. The ultimate behavior of such a sequence has already been studied in detail by several authors if the matrix is irreducible (see [1, 5, 7, 8] and the references therein). For reducible matrices it has been shown that the ultimate behavior is periodic [8, 9]. We will extend these results (and correct the results of [7] and [13]) by completely characterizing the rates and periods of the ultimate behavior of the entries of the sequence of consecutive powers of a general max-plus-algebraic matrix.

Our main motivation for studying this problem lies in the max-plus-algebraic system theory for discrete event systems. Typical examples of discrete event systems are flexible manufacturing systems, telecommunication networks, parallel processing systems, traffic control systems and logistic systems. The class of discrete event systems essentially consists of man-made systems that contain a finite number of resources (e.g. machines, communications channels, or processors) that are shared by several users (e.g. product types, information packets, or jobs) all of which contribute to the achievement of some common goal (e.g. the assembly of products, the end-to-end transmission of a set of information packets, or a parallel computation) [1]. There are many modeling techniques for discrete event systems, such as (extended) state machines, max-plus algebra, formal languages, automata, temporal logic, generalized semi-Markov processes, Petri nets, computer simulation models and so on (see [1, 4, 12, 11] and the references cited therein). In general models that describe the behavior of a discrete event system are nonlinear in conventional algebra. However, there is a class

*Control Laboratory, Faculty of Information Technology and Systems, Delft University of Technology, P.O. Box 5031, 2600 GA Delft, The Netherlands, email: b.deschutter@its.tudelft.nl

of discrete event systems – the max-plus-linear discrete event systems – that can be described by a model that is “linear” in the max-plus algebra [1, 5, 6]. The model of a max-plus-linear discrete event system can be characterized by a triple of matrices (A, B, C) , which are called the system matrices of the model. The ultimate behavior of the system matrix A determines the ultimate behavior of the max-plus-linear discrete event system [1, 6].

This paper is organized as follows. In Section 2 we introduce some notation and we give a short introduction to the max-plus algebra and to graph theory. We also discuss the connection between max-plus-algebraic matrix operations and graph theory. In Section 3 we characterize the rates and periods of the ultimate behavior of the entries of the sequence of consecutive powers of a general max-plus-algebraic matrix. Finally we present some conclusions in Section 4.

2 Notation and definitions

2.1 Notation

The set of the real numbers is denoted by \mathbb{R} , the set of the nonnegative integers by \mathbb{N} , and the set of the positive integers by \mathbb{N}_0 . The number of elements of a set γ is denoted by $\#\gamma$. The least common multiple of the elements of a set γ of positive integers is denoted by $\text{lcm } \gamma$.

Let $A \in \mathbb{R}^{m \times n}$. The entry on the i th row and the j th column of A is denoted by a_{ij} or $(A)_{ij}$. If $\alpha \subseteq \{1, \dots, m\}$ and $\beta \subseteq \{1, \dots, n\}$, then $A_{\alpha\beta}$ is the submatrix of A obtained by removing all rows not indexed by α and all columns not indexed by β .

2.2 Max-plus algebra

The basic operations of the max-plus algebra are the maximum (represented by \oplus) and the addition (represented by \otimes):

$$\begin{aligned} x \oplus y &= \max(x, y) \\ x \otimes y &= x + y \end{aligned}$$

with $x, y \in \mathbb{R} \cup \{-\infty\}$. Define $\varepsilon = -\infty$ and $\mathbb{R}_\varepsilon = \mathbb{R} \cup \{\varepsilon\}$. The operations \oplus and \otimes are extended to matrices as follows. If $A, B \in \mathbb{R}_\varepsilon^{m \times n}$ and $C \in \mathbb{R}_\varepsilon^{n \times p}$ then we have

$$\begin{aligned} (A \oplus B)_{ij} &= a_{ij} \oplus b_{ij} \\ (A \otimes C)_{ij} &= \bigoplus_{k=1}^p a_{ik} \otimes c_{kj} \end{aligned}$$

for all i, j . Note that these definitions resemble the definitions of the sum and the product of matrices in linear algebra but with \oplus instead of $+$ and \otimes instead of \times . This analogy is one of the reasons why we call \oplus the max-plus-algebraic addition and \otimes the max-plus-algebraic multiplication. For more information on the analogies and differences between max-plus algebra and linear algebra the interested reader is referred to [1, 6].

The matrix $\mathcal{E}_{m \times n}$ is the $m \times n$ max-plus-algebraic zero matrix: $(\mathcal{E}_{m \times n})_{ij} = \varepsilon$ for all i, j . The matrix E_n is the $n \times n$ max-plus-algebraic identity matrix: $(E_n)_{ii} = 0$ for all i and $(E_n)_{ij} = \varepsilon$ for all i, j with $i \neq j$. If we permute the rows or the columns of E_n , we obtain a max-plus-algebraic permutation matrix. For a max-plus-algebraic permutation matrix $P \in \mathbb{R}_\varepsilon^{n \times n}$ we have $P \otimes P^T = P^T \otimes P = E_n$.

Let $k \in \mathbb{N}$. The k th max-plus-algebraic power of $x \in \mathbb{R}$ is denoted by $x^{\otimes k}$ and corresponds to kx in conventional algebra. If $k > 0$ then $\varepsilon^{\otimes k} = \varepsilon$. We have $\varepsilon^{\otimes 0} = 0$ by definition. The max-plus-algebraic matrix power of $A \in \mathbb{R}_\varepsilon^{n \times n}$ is defined as follows:

$$A^{\otimes 0} = E_n \quad \text{and} \quad A^{\otimes k} = A \otimes A^{\otimes k-1} \quad \text{for } k = 1, 2, \dots$$

Consider sequences $h_i = \{(h_i)_k\}_{k=0}^\infty$ for $i = 1, \dots, m$ with $(h_i)_k \in \mathbb{R}_\varepsilon$ for all i, k . The max-plus-algebraic sum $g = h_1 \oplus \dots \oplus h_m$ is defined by $g_k = (h_1)_k \oplus \dots \oplus (h_m)_k$. The max-plus-algebraic product $g = h_1 \otimes \dots \otimes h_m$ is defined by

$$g_k = \bigoplus_{\substack{k_1, \dots, k_m \in \mathbb{N} \\ k_1 + \dots + k_m = k}} (h_1)_{k_1} \otimes \dots \otimes (h_m)_{k_m} .$$

Definition 2.1 (Ultimately geometric sequence) *We say that the sequence $\{g_k\}_{k=0}^\infty$ is ultimately geometric if*

$$\exists K \in \mathbb{N}, \exists c \in \mathbb{N}_0, \exists \lambda \in \mathbb{R}_\varepsilon \text{ such that } \forall k \geq K : g_{k+c} = \lambda^{\otimes c} \otimes g_k . \quad (1)$$

The term ‘‘ultimately geometric’’ was introduced by Gaubert in [8, 9]. Note that ‘‘geometric’’ has to be understood in the max-plus-algebraic sense: the terms of the sequence are max-plus-multiplied by a constant factor $c\lambda$. If g is an ultimately geometric sequence then the smallest possible c for which (1) holds is called the *period* of g . The smallest possible corresponding λ is then called the *rate* of g . Note that $\{\varepsilon\}_{k=0}^\infty$ has period 1 and rate ε .

Definition 2.2 (Ultimately periodic sequence) *We say that the sequence $\{g_k\}_{k=0}^\infty$ is ultimately periodic if*

$$\begin{aligned} \exists K \in \mathbb{N}, \exists c \in \mathbb{N}_0, \exists \lambda_0, \dots, \lambda_{c-1} \in \mathbb{R}_\varepsilon \text{ such that} \\ g_{kc+c+s} = \lambda_s^{\otimes c} \otimes g_{kc+s} \text{ for all } k \geq K \text{ and for } s = 0, \dots, c-1. \end{aligned} \quad (2)$$

If g is an ultimately periodic sequence then the smallest possible c for which (2) holds is called the *period* of g . The smallest possible corresponding λ_s s are called the *rates* of g . In general the max-plus-algebraic sum of ultimately geometric sequences is ultimately periodic. The reverse also holds: every ultimately periodic sequence can be considered as the max-plus-algebraic sum of ultimately geometric sequences [8, 9].

Let us now illustrate the concepts defined above by an example.

Example 2.3 Consider the sequence

$$g = \{g_k\}_{k=0}^\infty = 0, 0, 0, 2, \varepsilon, 0, 8, \varepsilon, 6, 14, \varepsilon, 12, 20, \varepsilon, 18, \dots$$

This sequence is ultimately geometric with rate $\lambda = 2$ and period $c = 3$ since $g_{k+3} = 2^{\otimes 3} \otimes g_k = 6 \otimes g_k = 6 + g_k$ for all $k \geq 3$. The sequence

$$h = \{h_k\}_{k=0}^\infty = \varepsilon, 0, 1, 0, 2, 0, 3, 0, 4, \dots$$

is ultimately periodic with period $c = 2$ and rates $\lambda_0 = \frac{1}{2}$ and $\lambda_1 = 0$ since $h_{2k+2} = \left(\frac{1}{2}\right)^{\otimes 2} \otimes h_{2k} = 1 \otimes h_{2k}$ and $h_{2k+2+1} = 0^{\otimes 2} \otimes h_{2k+1} = h_{2k+1}$ for all $k \geq 1$.

It is easy to verify that h can be written as the max-plus-algebraic sum of the ultimately geometric sequences $h_1 = \varepsilon, \varepsilon, 1, \varepsilon, 2, \varepsilon, 3, \varepsilon, 4, \dots$ and $h_2 = \varepsilon, 0, \varepsilon, 0, \varepsilon, 0, \varepsilon, 0, \dots$. Note that the rates of h_1 and h_2 are respectively $\frac{1}{2}$ and 0, and that their period is equal to 2. \square

2.3 Max-plus-algebra and graph theory

We assume that the reader is familiar with basic concepts of graph theory such as directed graph, loop, circuit, elementary circuit and so on (see e.g. [1, 14]).

If we have a directed graph \mathcal{G} with vertex set $\mathcal{V} = \{1, 2, \dots, n\}$ and if we associate a real number w_{ij} with each arc (j, i) of \mathcal{G} , then we say that \mathcal{G} is a weighted directed graph. We call w_{ij} the weight of the arc (j, i) . Note that the first subscript of w_{ij} corresponds to the final (and not the initial) vertex of the arc (j, i) . With every weighted directed graph \mathcal{G} with vertex set $\mathcal{V} = \{1, 2, \dots, n\}$ there corresponds a matrix $A \in \mathbb{R}_\varepsilon^{n \times n}$ such that $a_{ij} = w_{ij}$ if there is an arc (j, i) in \mathcal{G} with weight w_{ij} and $a_{ij} = \varepsilon$ if there is no arc (j, i) in \mathcal{G} . We say that \mathcal{G} is the precedence graph of A , denoted by $\mathcal{G}(A)$.

Let $A \in \mathbb{R}_\varepsilon^{n \times n}$. The weight of a path $i_1 \rightarrow i_2 \rightarrow \dots \rightarrow i_l$ in $\mathcal{G}(A)$ is defined as the sum of the weights of the arcs that compose the path. Let us now give a graph-theoretic interpretation of the max-plus-algebraic matrix power. If $k \in \mathbb{N}_0$ then we have

$$\begin{aligned} (A^{\otimes k})_{ij} &= \bigoplus_{i_1, \dots, i_{k-1}} a_{ii_1} \otimes a_{i_1 i_2} \otimes \dots \otimes a_{i_{k-1} j} \\ &= \max_{i_1, \dots, i_{k-1}} (a_{ii_1} + a_{i_1 i_2} + \dots + a_{i_{k-1} j}) \end{aligned}$$

for all i, j . Hence, $(A^{\otimes k})_{ij}$ is the maximal weight of all paths of $\mathcal{G}(A)$ of length k that have j as their initial vertex and i as their final vertex — where the maximal weight is equal to ε by definition if there does not exist a path of length k from j to i .

The average weight of a circuit is defined as the weight of the circuit divided by its length. A circuit is called critical if it has maximum average weight. The critical graph $\mathcal{G}^c(A)$ of A consists of those vertices and arcs of $\mathcal{G}(A)$ that belong to some critical circuit of $\mathcal{G}(A)$.

We say that $\mathcal{G}(A)$ is strongly connected if for any two *different*¹ vertices v_i, v_j of $\mathcal{G}(A)$ there exists a path from v_i to v_j . A maximal strongly connected subgraph (m.s.c.s.) \mathcal{G}_{sub} of $\mathcal{G}(A)$ is a strongly connected subgraph that is maximal, i.e. if we add an extra vertex (and some extra arcs) of $\mathcal{G}(A)$ to \mathcal{G}_{sub} then \mathcal{G}_{sub} is no longer strongly connected. The matrix A is called irreducible if $\mathcal{G}(A)$ is strongly connected.

The cyclicity of an irreducible matrix A is equal to the greatest common divisor of the lengths of all the elementary circuits of the $\mathcal{G}^c(A)$. If the graph $\mathcal{G}^c(A)$ contains no circuits then the cyclicity is equal to 1 by definition. Note that the 1×1 max-plus-algebraic zero matrix $[\varepsilon]$ is the only max-plus-algebraic zero matrix that is irreducible and that its cyclicity is equal to 1. The following theorem gives a relation between the cyclicity of an irreducible matrix A and the ultimate behavior of the sequence $\{A^{\otimes k}\}_{k=0}^\infty$.

Theorem 2.4 *If $A \in \mathbb{R}_\varepsilon^{n \times n}$ is irreducible, then*

$$\exists \lambda \in \mathbb{R}_\varepsilon, \exists k_0 \in \mathbb{N} \text{ such that } \forall k \geq k_0 : A^{\otimes k+c} = \lambda^{\otimes c} \otimes A^{\otimes k}$$

¹Most authors do not add the extra condition that the vertices should be different. However, this definition, which was taken from [1], makes some of the subsequent definitions, theorems and proofs easier to formulate.

where c is the cyclicity of A .

Proof: See e.g. Theorem 1.2.3 of [8]. ■

The number λ that appears in Theorem 2.4 is called the max-plus-algebraic eigenvalue of A and it corresponds to the maximal average weight over all elementary circuits of $\mathcal{G}(A)$.

The following theorem is the max-plus-algebraic analogue of a well-known result from matrix algebra that states that any square matrix can be transformed into a block upper diagonal matrix with irreducible blocks by simultaneously reordering the rows and columns of the matrix (see e.g. [1, 3, 10] for the proof of this theorem and for its interpretation in terms of graph theory):

Theorem 2.5 *If $A \in \mathbb{R}_\varepsilon^{n \times n}$ then there exists a max-plus-algebraic permutation matrix $P \in \mathbb{R}_\varepsilon^{n \times n}$ such that the matrix $\hat{A} = P \otimes A \otimes P^T$ is a max-plus-algebraic block upper triangular matrix of the form*

$$\hat{A} = \begin{bmatrix} \hat{A}_{11} & \hat{A}_{12} & \dots & \hat{A}_{1l} \\ \varepsilon & \hat{A}_{22} & \dots & \hat{A}_{2l} \\ \vdots & \vdots & \ddots & \vdots \\ \varepsilon & \varepsilon & \dots & \hat{A}_{ll} \end{bmatrix} \quad (3)$$

with $l \geq 1$ and where the matrices $\hat{A}_{11}, \dots, \hat{A}_{ll}$ are square and irreducible. The matrices $\hat{A}_{11}, \dots, \hat{A}_{ll}$ are uniquely determined to within simultaneous permutation of their rows and columns, but their ordering in (3) is not necessarily unique.

The form in (3) is called the max-plus-algebraic Frobenius normal form of A . If A is irreducible then there is only one block in (3) and then A is a max-plus-algebraic Frobenius normal form of itself. Each diagonal block of \hat{A} corresponds to an m.s.c.s. of $\mathcal{G}(\hat{A})$. If $\hat{A} = P \otimes A \otimes P^T$ is the max-plus-algebraic Frobenius normal form of A , then we have $A = P^T \otimes \hat{A} \otimes P$ since P is a max-plus-algebraic permutation matrix. Hence,

$$A^{\otimes k} = (P^T \otimes \hat{A} \otimes P)^{\otimes k} = P^T \otimes \hat{A}^{\otimes k} \otimes P \quad \text{for all } k \in \mathbb{N}. \quad (4)$$

3 The ultimate behavior of the sequence $\{A^{\otimes k}\}_{k=0}^\infty$

If $A \in \mathbb{R}_\varepsilon^{n \times n}$ is irreducible the ultimate behavior of $\{A^{\otimes k}\}_{k=0}^\infty$ is characterized by Theorem 2.4. For a general matrix A it has already been shown in [8] that the sequences $\{(A^{\otimes k})_{ij}\}_{k=0}^\infty$ are ultimately periodic. An essentially equivalent result has been obtained independently by Bonnier-Rigny and Krob in [2] for the structure $(\mathbb{N} \cup \{+\infty\}, \min, +)$, which is called the tropical semiring and which is strongly related to the max-plus-algebra. Now we will extend these results and give a detailed characterization of rates and periods of the entries of the sequence $\{A^{\otimes k}\}_{k=0}^\infty$ for a general matrix A .

The following two technical lemmas will be used in the proof of the main theorem. Their proofs can be found in the appendix.

Lemma 3.1 Consider m ultimately geometric sequences h_1, \dots, h_m with rates different from ε . Let c_i be the period of h_i and let λ_i be the rate of h_i for $i = 1, \dots, m$. If $g = h_1 \oplus \dots \oplus h_m$ and if $c = \text{lcm}(c_1, \dots, c_m)$ then

$$\begin{aligned} \exists K \in \mathbb{N}, \exists \gamma_0, \dots, \gamma_{c-1} \in \{\lambda_1, \dots, \lambda_m\} \text{ such that} \\ g_{kc+c+s} = \gamma_s^{\otimes c} \otimes g_{kc+s} \quad \text{for all } k \geq K \text{ and for } s = 0, \dots, c-1. \end{aligned} \quad (5)$$

Furthermore, there exists at least one index $s \in \{0, \dots, c-1\}$ such that the smallest γ_s for which (5) holds is equal to $\bigoplus_{i=1}^m \lambda_i$.

Lemma 3.2 Consider m ultimately geometric sequences h_1, \dots, h_m with rates different from ε . Let c_i be the period of h_i and let λ_i be the rate of h_i for $i = 1, \dots, m$. If $g = h_1 \otimes \dots \otimes h_m$ and if $c = \text{lcm}(c_1, \dots, c_m)$ then

$$\begin{aligned} \exists K \in \mathbb{N}, \exists \gamma_0, \dots, \gamma_{c-1} \in \mathbb{R}_\varepsilon \text{ such that} \\ g_{kc+c+s} = \gamma_s^{\otimes c} \otimes g_{kc+s} \quad \text{for all } k \geq K \text{ and for } s = 0, \dots, c-1. \end{aligned} \quad (6)$$

There exists at least one index $s \in \{0, 1, \dots, c-1\}$ such that the smallest γ_s for which (6) holds is equal to $\bigoplus_{i=1}^m \lambda_i$. Moreover, for k^* large enough $\{g_k\}_{k=k^*}^\infty$ can be written as a finite sum of ultimately geometric sequences with rates λ_i and periods c_i .

If \hat{A} is a max-plus-algebraic normal form of A , then it follows from (4) that we may consider the sequence $\{\hat{A}^{\otimes k}\}_{k=0}^\infty$ instead of $\{A^{\otimes k}\}_{k=0}^\infty$ if we want to study the ultimate behavior of the sequence of consecutive powers of A . The following theorem, which is an extension of Theorem 2.4 and a corrected version of Lemma 4 of [13] and of Lemma C.1.4 of [7] characterizes the rates and periods of the ultimate behavior of $\{\hat{A}^{\otimes k}\}_{k=0}^\infty$:

Theorem 3.3 Let $\hat{A} \in \mathbb{R}_\varepsilon^{n \times n}$ be a matrix of the form (3) where the matrices $\hat{A}_{11}, \dots, \hat{A}_{ll}$ are square and irreducible. Let λ_i and c_i be respectively the max-plus-algebraic eigenvalue and the cyclicity of \hat{A}_{ii} for $i = 1, \dots, l$. Define sets $\alpha_1, \dots, \alpha_l$ such that $\hat{A}_{\alpha_i \alpha_j} = \hat{A}_{ij}$ for all i, j with $i \leq j$.

Define

$$S_{ij} = \{ \{i_0, \dots, i_s\} \subseteq \{1, \dots, l\} \mid i = i_0 < i_1 < \dots < i_s = j \text{ and}$$

$$\hat{A}_{i_r i_{r+1}} \neq \varepsilon \text{ for } r = 0, \dots, s-1 \}$$

$$\Gamma_{ij} = \bigcup_{\gamma \in S_{ij}} \gamma$$

$$\Lambda_{ij} = \begin{cases} \{\lambda_t \mid t \in \Gamma_{ij}\} & \text{if } \Gamma_{ij} \neq \emptyset, \\ \{\varepsilon\} & \text{if } \Gamma_{ij} = \emptyset, \end{cases}$$

$$c_{ij} = \begin{cases} \text{lcm}\{c_t \mid t \in \Gamma_{ij}\} & \text{if } \Gamma_{ij} \neq \emptyset \text{ and } c_t \neq 0 \text{ for some } t \in \Gamma_{ij}, \\ 1 & \text{otherwise,} \end{cases}$$

for all i, j with $i < j$. We have

$$\forall i, j \in \{1, \dots, l\} \text{ with } i > j : \left(\hat{A}^{\otimes k} \right)_{\alpha_i \alpha_j} = \mathcal{E}_{n_i \times n_j} \quad \text{for all } k \in \mathbb{N}. \quad (7)$$

Moreover, there exists an integer $K \in \mathbb{N}$ such that

$$\forall i \in \{1, \dots, l\} : \left(\hat{A}^{\otimes k+c_i} \right)_{\alpha_i \alpha_i} = \lambda_i^{\otimes c_i} \otimes \left(\hat{A}^{\otimes k} \right)_{\alpha_i \alpha_i} \quad \text{for all } k \geq K \quad (8)$$

and

$\forall i, j \in \{1, \dots, l\}$ with $i < j, \forall p \in \alpha_i, \forall q \in \alpha_j, \exists \gamma_0, \dots, \gamma_{c_{ij}-1} \in \Lambda_{ij}$ such that

$$\left(\hat{A}^{\otimes kc_{ij}+c_{ij}+s} \right)_{pq} = \gamma_s^{\otimes c_{ij}} \otimes \left(\hat{A}^{\otimes kc_{ij}+s} \right)_{pq} \quad \text{for all } k \geq K \text{ and for } s = 0, \dots, c_{ij} - 1. \quad (9)$$

Furthermore, for each combination i, j, p, q with $i < j, p \in \alpha_i$ and $q \in \alpha_j$, there exists at least one index $s \in \{0, \dots, c_{ij} - 1\}$ such that the smallest γ_s for which (9) holds is equal to $\max \Lambda_{ij}$.

Remark 3.4 Let us give a graphical interpretation of the sets S_{ij} and Γ_{ij} . Let C_i be the m.s.c.s. of $\mathcal{G}(\hat{A})$ that corresponds to \hat{A}_{ii} for $i = 1, \dots, l$. So α_i is the vertex set of C_i .

If $\{i_0 = i, i_1, \dots, i_s = j\} \in S_{ij}$ then there exists a path from a vertex in C_{i_r} to a vertex in $C_{i_{r-1}}$ for each $r \in \{1, \dots, s\}$. Since each m.s.c.s. C_i of $\mathcal{G}(\hat{A})$ is strongly connected, this implies that there exists a path from a vertex in C_j to a vertex in C_i that passes through $C_{i_{s-1}}, C_{i_{s-2}}, \dots, C_{i_1}$.

If $S_{ij} = \emptyset$ then there does not exist any path from a vertex in C_j to a vertex in C_i .

The set Γ_{ij} is the set of indices of the m.s.c.s.'s of $\mathcal{G}(\hat{A})$ through which some path from a vertex of C_j to a vertex of C_i passes. \diamond

Proof of Theorem 3.3: Since the matrices $\hat{A}_{\alpha_i \alpha_i}$ are irreducible (8) is a direct consequence of Theorem 2.4.

Recall that $(\hat{A}^{\otimes k})_{ij}$ is equal to the maximal weight over all paths of length k from j to i in $\mathcal{G}(\hat{A})$ where the maximal weight is equal to ε by definition if there does not exist any path of length k from j to i . Let C_i be the m.s.c.s. of $\mathcal{G}(\hat{A})$ that corresponds to \hat{A}_{ii} for $i = 1, \dots, l$. Since $\hat{A}_{\alpha_i \alpha_j} = \mathcal{E}_{n_i \times n_j}$ if $i > j$, there are no arcs from any vertex of C_j to a vertex in C_i . As a consequence, (7) holds.

Now consider $i, j \in \{1, \dots, l\}$ with $i < j$. We distinguish three cases:

- If $\Gamma_{ij} = \emptyset$ then there does not exist a path from a vertex in C_j to a vertex in C_i . Hence, $(\hat{A}^{\otimes k})_{\alpha_i \alpha_j} = \mathcal{E}_{n_i \times n_j}$ for all $k \in \mathbb{N}$. Since in this case we have $\Lambda_{ij} = \{\varepsilon\}$ and $c_{ij} = 1$, this implies that (9) and the last statement of the theorem hold if $\Gamma_{ij} = \emptyset$.
- If $\Gamma_{ij} \neq \emptyset$ and $\Lambda_{ij} = \{\varepsilon\}$ then $\hat{A}_{tt} = [\varepsilon]$ and $c_t = 1$ for all $t \in \Gamma_{ij}$. So there exist paths from a vertex in C_j to a vertex in C_i , but each path passes only through m.s.c.s.'s that consist of one vertex and contain no loop. Such a path passes through at most $\#\Gamma_{ij}$ of such m.s.c.s.'s (C_j and C_i included). This implies that there does not exist a path with a length larger than or equal to $\#\Gamma_{ij}$ from a vertex in C_j to a vertex in C_i . Hence, $(\hat{A}^{\otimes k})_{\alpha_i \alpha_j} = \mathcal{E}_{n_i \times n_j}$ for all $k \geq \#\Gamma_{ij}$. Furthermore, $c_{ij} = 1$ since $c_t = 1$ for all $t \in \Gamma_{ij}$. Hence, (9) and the last statement of the theorem also hold if $\Gamma_{ij} = \emptyset$ and $\Lambda_{ij} = \{\varepsilon\}$.




Figure 1: Illustration of the proof of Theorem 3.3. There exists a path from vertex u_s of m.s.c.s. C_j to vertex v_0 of m.s.c.s. C_i that passes through the m.s.c.s.'s $C_{i_{s-1}}, C_{i_{s-2}}, \dots, C_{i_1}$.

- Finally, we consider the case with $\Gamma_{ij} \neq \emptyset$ and $\Lambda_{ij} \neq \{\varepsilon\}$. Select an arbitrary vertex p of C_i and an arbitrary vertex q of C_j . For each set $\gamma = \{i_0, \dots, i_s\} \in S_{ij}$ we define

$$\mathcal{S}(\gamma) = \{(U, V) \mid U = \{u_0, \dots, u_s\}, V = \{v_0, \dots, v_s\}, u_s = q, v_0 = p, \text{ and} \\ u_r \in \alpha_{i_r}, v_{r+1} \in \alpha_{i_{r+1}} \text{ and } (\hat{A})_{u_r v_{r+1}} \neq \varepsilon \text{ for } r = 0, \dots, s\} .$$

So if $(U, V) \in \mathcal{S}(\gamma)$ with $U = \{u_0, \dots, u_s\}$ and $V = \{v_0, \dots, v_s\}$ then there exists a path from q to p that passes through m.s.c.s. C_{i_r} for $r = 0, \dots, s$ and that enters C_{i_r} at vertex u_r for $r = 0, \dots, s-1$ and that exits from C_{i_r} through vertex v_r for $r = 1, \dots, s$ (see also Figure 1). Hence, we have

$$\left(\hat{A}^{\otimes k}\right)_{pq} = \bigoplus_{\gamma \in S_{ij}} \bigoplus_{(U, V) \in \mathcal{S}(\gamma)} g(\gamma, U, V) \quad \text{for all } k \in \mathbb{N}_0$$

where

$$g(\gamma, U, V) = \bigoplus_{\substack{p_0, \dots, p_s \in \mathbb{N} \\ p_0 + \dots + p_s = k-s}} (\hat{A}^{\otimes p_0})_{p u_0} \otimes (\hat{A}^{\otimes p_1})_{u_0 v_1} \otimes \dots \\ \otimes (\hat{A}^{\otimes p_{s-1}})_{u_{s-1} v_s} \otimes (\hat{A}^{\otimes p_s})_{v_s q} \quad (10)$$

with the empty max-plus-algebraic sum equal to ε by definition. Each term of the max-plus-algebraic sum in (10) represents the maximal weight over all paths from q to p that consist of the concatenation of paths of length p_r from vertex u_r to vertex v_r of

C_{i_r} for $r = 0, \dots, s$ and paths of length 1 from vertex v_{r+1} of $C_{i_{r+1}}$ to vertex u_r of C_{i_r} for $r = 0, \dots, s$ where by definition the maximal weight is equal to ε if no such paths exist. Note that if $\lambda_{i_r} = \varepsilon$ for some r then every term in the max-plus-algebraic sum (10) for which $p_r > 0$ will be equal to ε . Furthermore, since $\varepsilon^{\otimes 0} = 0$ by definition, this means that each factor of the form $(\hat{A}_{i_r i_r}^{\otimes p_r})_{u_r v_r}$ for which $\lambda_{i_r} = \varepsilon$ may be removed from the max-plus-algebraic sum (10). Note that indices t for which $\lambda_t = \varepsilon$ or equivalently $c_t = 1$ do not influence the value of c_{ij} . Also note that since $\Gamma_{ij} \neq \emptyset$ and $\Lambda_{ij} \neq \{\varepsilon\}$ we have at least one combination γ, U, V for which the sequence (10) has a rate λ_{i_r} that is different from ε .

Since $\hat{A}_{i_r i_r}$ is irreducible, we have

$$\left(\hat{A}_{i_r i_r}^{\otimes k+c_{i_r}}\right)_{v_r u_r} = \lambda_{i_r}^{\otimes c_{i_r}} \otimes \left(\hat{A}_{i_r i_r}^{\otimes k}\right)_{v_r u_r} \quad \text{for } k \text{ large enough}$$

by Theorem 2.4. Hence, if $g(\gamma, U, V)$ is different from ε , i.e. if it still contains terms after the factors for which $\lambda_{i_r} = \varepsilon$ have been removed, $g(\gamma, U, V)$ is a max-plus-algebraic product of ultimate geometric sequences with rates $\lambda_{i_r} \neq \varepsilon$ and periods c_{i_r} . From Lemma 3.2 it follows that $g(\gamma, U, V)$ is an ultimately periodic sequence and that for k^* large enough $\{(g(\gamma, U, V))_k\}_{k=k^*}^{\infty}$ can be written as the max-plus-algebraic sum of a finite number of ultimately geometric sequences with rates $\lambda_{i_r} \neq \varepsilon$ and periods c_{i_r} . So $\{(\hat{A}^{\otimes k})_{pq}\}_{k=0}^{\infty}$ is a max-plus-algebraic sum of ultimately geometric sequences with rates $\lambda_{i_r} \neq \varepsilon$ and periods c_{i_r} . Hence, it follows from Lemma 3.1 that (9) and the last statement of the theorem hold. \blacksquare

The following example shows that the lcm in the definition of c_{ij} in Theorem 3.3 is necessary (Lemma 4 of [13] incorrectly uses max instead of lcm.).

Example 3.5 Consider the matrix

$$A = \begin{array}{c|ccc|ccc|c} \varepsilon & 0 & \varepsilon & 0 & \varepsilon & \varepsilon & \varepsilon & \varepsilon \\ \varepsilon & \varepsilon & 0 & \varepsilon & \varepsilon & \varepsilon & 0 & 0 \\ \varepsilon & 0 & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon \\ \hline \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & 0 & 0 & 0 \\ \varepsilon & \varepsilon & \varepsilon & 0 & \varepsilon & \varepsilon & \varepsilon & \varepsilon \\ \varepsilon & \varepsilon & \varepsilon & \varepsilon & 0 & \varepsilon & \varepsilon & \varepsilon \\ \hline \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon & \varepsilon \end{array} .$$

This matrix is in max-plus-algebraic Frobenius normal form and its block structure is indicated by the vertical and horizontal lines. The precedence graph of A is represented in Figure 2. The sets and variables of Theorem 3.3 have the following values for A : $\alpha_1 = \{1\}$, $\alpha_2 = \{2, 3\}$, $\alpha_3 = \{4, 5, 6\}$, $\alpha_4 = \{7\}$, $\lambda_1 = \lambda_4 = \varepsilon$, $\lambda_2 = \lambda_3 = 0$, $c_1 = c_4 = 1$, $c_2 = 2$ and $c_3 = 3$. Now we consider the ultimate behavior of the sequence $\{(A^{\otimes k})_{\alpha_1 \alpha_4}\}_{k=0}^{\infty}$. Note that $S_{14} = \{\{2\}, \{3\}\}$, $\Gamma_{14} = \{2, 3\}$, $\Gamma_{23} = \{0\}$, and $c_{14} = \text{lcm}(c_2, c_3) = \text{lcm}(2, 3) = 6$. We have

$$\{(A^{\otimes k})_{\alpha_1 \alpha_4}\}_{k=0}^{\infty} = \varepsilon, 0, \varepsilon, 0, 0, 0, \varepsilon, 0, \varepsilon, 0, 0, 0, \varepsilon, 0, \varepsilon, 0, 0, 0, \varepsilon, 0, \varepsilon, 0, 0, 0, \dots$$

The period of this sequence is given by $c_{14} = 6 = \text{lcm}(c_2, c_3)$. Hence, the lcm in the definition of c_{ij} in Theorem 3.3 is really necessary. \square




Figure 2: The precedence graph $\mathcal{G}(A)$ of the matrix A of Example 3.5. All the arcs have weight 0.

The following example shows that the sequence $\{(\hat{A}^{\otimes k})_{ij}\}_{k=1}^{\infty}$ is in general not ultimately geometric (Lemma 4 of [13] and Lemma C.1.4 of [7] incorrectly state that if $i < j$ then the matrix sequence $\{\hat{A}^{\otimes k}\}_{k=0}^{\infty}$ is ultimately geometric).

Example 3.6 We construct the matrix \tilde{A} from the matrix A of Example 3.5 by replacing a_{23} by 2 and keeping all other entries. Now we have $\lambda_2 = 1$. The values of the other variables and sets of Theorem 3.3 are the same as for the matrix A of Example 3.5. We have

$$\{(\tilde{A}^{\otimes k})_{\alpha_1 \alpha_4}\}_{k=0}^{\infty} = \varepsilon, 0, \varepsilon, 2, 0, 4, \varepsilon, 6, \varepsilon, 8, 0, 10, \varepsilon, 12, \varepsilon, 14, 0, 16, \varepsilon, 18, \varepsilon, 20, 0, 22, \dots$$

This sequence is ultimately periodic with period $c_{14} = 6$ and with rates $\gamma_0 = \gamma_2 = \varepsilon$, $\gamma_1 = \gamma_3 = \gamma_5 = 1 = \lambda_2$ and $\gamma_4 = 0 = \lambda_4$. So the sequence $\{(\hat{A}^{\otimes k})_{ij}\}_{k=0}^{\infty}$ is in general not ultimately geometric. \square

Corollary 3.7 Let $\hat{A} \in \mathbb{R}_{\varepsilon}^{n \times n}$ be a matrix of the form (3) where the matrices $\hat{A}_{11}, \hat{A}_{22}, \dots, \hat{A}_{ll}$ are square and irreducible. Let λ_i and c_i be respectively the max-plus-algebraic eigenvalue and the cyclicity of \hat{A}_{ii} for $i = 1, \dots, l$. Let α_i, Λ_{ij} and c_{ij} be defined as in Theorem 3.3. Then there exists an integer K such that

$$\begin{aligned} \forall i, j \in \{1, \dots, l\} \text{ with } i > j, \forall p \in \alpha_i, \forall q \in \alpha_j, \exists s \in \{0, \dots, c_{ij} - 1\} \text{ such that} \\ \left(\hat{A}^{\otimes kc_{ij} + s + c_{ij}} \oplus \hat{A}^{\otimes kc_{ij} + s + c_{ij} + 1} \oplus \dots \oplus \hat{A}^{\otimes kc_{ij} + s + 2c_{ij} - 1} \right)_{pq} = \\ \lambda_{ij}^{\otimes c_{ij}} \otimes \left(\hat{A}^{\otimes kc_{ij} + s} \oplus \hat{A}^{\otimes kc_{ij} + s + 1} \oplus \dots \oplus \hat{A}^{\otimes kc_{ij} + s + c_{ij} - 1} \right)_{pq} \quad \text{for all } k \geq K, \end{aligned}$$

where $\lambda_{ij} = \max \Lambda_{ij}$.

Proof: This is a direct consequence of the last statement of Theorem 3.3. \blacksquare

4 Conclusions

In this paper we have given a detailed characterization of the rates and periods of the ultimate behavior of the sequence of consecutive matrix powers of a general max-plus-algebraic matrix as a function of structural parameters of the matrix such as its size, its Frobenius normal form, and the eigenvalues and cyclicities of the diagonal blocks in the Frobenius normal form. This result extends and corrects previously known results. We have only considered the ultimate behavior of the sequence. An important open question and topic for future research is the characterization of both the length and the evolution of the *transient* behavior of the sequence as a function of the structural parameters of the matrix.

Acknowledgment

We would like to thank the anonymous referees for their careful review and useful comments, which certainly contributed to improving the clarity of this paper, and for pointing out an error in the original proof of Theorem 3.3.

References

- [1] F. Baccelli, G. Cohen, G.J. Olsder, and J.P. Quadrat, *Synchronization and Linearity*. New York: John Wiley & Sons, 1992.
- [2] A. Bonnier-Rigny and D. Krob, "A complete system of identities for one-letter rational expressions with multiplicities in the tropical semiring," *Theoretical Computer Science*, vol. 134, no. 1, pp. 27–50, Nov. 1994.
- [3] R.A. Brualdi and H.J. Ryser, *Combinatorial Matrix Theory*, vol. 39 of *Encyclopedia of Mathematics and Its Applications*. Cambridge, UK: Cambridge University Press, 1991.
- [4] C.G. Cassandras, S. Lafortune, and G.J. Olsder, "Introduction to the modelling, control and optimization of discrete event systems," in *Trends in Control: A European Perspective* (A. Isidori, ed.), pp. 217–291, Berlin, Germany: Springer-Verlag, 1995.
- [5] G. Cohen, D. Dubois, J.P. Quadrat, and M. Viot, "A linear-system-theoretic view of discrete-event processes and its use for performance evaluation in manufacturing," *IEEE Transactions on Automatic Control*, vol. 30, no. 3, pp. 210–220, Mar. 1985.
- [6] R.A. Cuninghame-Green, *Minimax Algebra*, vol. 166 of *Lecture Notes in Economics and Mathematical Systems*. Berlin, Germany: Springer-Verlag, 1979.
- [7] B. De Schutter, *Max-Algebraic System Theory for Discrete Event Systems*. PhD thesis, Faculty of Applied Sciences, K.U.Leuven, Leuven, Belgium, Feb. 1996.
- [8] S. Gaubert, "On rational series in one variable over certain dioids," Tech. rep. 2162, INRIA, Le Chesnay, France, Jan. 1994.
- [9] S. Gaubert, "Rational series over dioids and discrete event systems," in *Proceedings of the 11th International Conference on Analysis and Optimization of Systems* (Sophia-Antipolis, France, June 1994) (G. Cohen and J.P. Quadrat, eds.), vol. 199 of *Lecture Notes in Control and Information Sciences*, pp. 247–256, Springer-Verlag: London, 1994.

- [10] F. Harary, "A graph theoretic approach to matrix inversion by partitioning," *Numerische Mathematik*, vol. 4, pp. 128–135, 1962.
- [11] Y.C. Ho, ed., *Proceedings of the IEEE*, vol. 77, no. 1, Jan. 1989. Special Issue on Dynamics of Discrete Event Systems.
- [12] Y.C. Ho, ed., *Discrete Event Dynamic Systems: Analyzing Complexity and Performance in the Modern World*. Piscataway, New Jersey: IEEE Press, 1992.
- [13] M. Wang, Y. Li, and H. Liu, "On periodicity analysis and eigen-problem of matrix in max-algebra," in *Proceedings of the 1991 IFAC Workshop on Discrete Event System Theory and Applications in Manufacturing and Social Phenomena*, Shenyang, China, pp. 44–48, June 1991.
- [14] R.J. Wilson, *Introduction to Graph Theory*. London: Longman Group, 2nd ed., 1979.

Appendix: Proofs of Lemmas 3.1 and 3.2

Proof of Lemma 3.1: In this proof we always assume that $i \in \{1, \dots, m\}$, $s, s^* \in \{0, \dots, c-1\}$ and $k, l \in \mathbb{N}$. Since each sequence h_i is ultimately geometric, there exists an integer K such that $(h_i)_{k+c_i} = \lambda_i^{\otimes c_i} \otimes (h_i)_k$ for all $k \geq K$ and for all i . Hence,

$$(h_i)_{k+pc_i} = \lambda_i^{\otimes pc_i} \otimes (h_i)_k \quad \text{for all } p \in \mathbb{N}, \text{ for all } k \geq K \text{ and for all } i. \quad (11)$$

Since $c = \text{lcm}(c_1, \dots, c_m)$ there exist positive integers w_1, \dots, w_m such that $c = w_i c_i$ for all i . Select $L \in \mathbb{N}$ with $Lc \geq K$. Consider an arbitrary index s . Since $Lc + s \geq K$, it follows from (11) that

$$(h_i)_{lc+s} = (h_i)_{Lc+s+(l-L)w_i c_i} = \lambda_i^{\otimes (l-L)w_i c_i} \otimes (h_i)_{Lc+s} = \lambda_i^{\otimes (l-L)c} \otimes (h_i)_{Lc+s} \quad (12)$$

for all $l \geq L$ and for all i . Define $N_s = \{i \mid (h_i)_{Lc+s} \neq \varepsilon\}$. We consider two cases:

- If $N_s = \emptyset$ then $(h_i)_{Lc+s} = \varepsilon$ for all i and thus also $(h_i)_{lc+s} = \varepsilon$ for all $l \geq L$ and for all i by (12). Hence, $g_{lc+s} = \varepsilon$ for all $l \geq L$. So if we set $\gamma_s = \lambda_1$ and select $K \geq K_s \stackrel{\text{def}}{=} L$ then (5) holds for this case.
- If $N_s \neq \emptyset$ then we define $\gamma_s = \max_{i \in N_s} \lambda_i$ and $i_s = \arg \max_{i \in N_s} \{(h_i)_{Lc+s} \mid \lambda_i = \gamma_s\}$. By (12) we have $(h_i)_{lc+s} = (h_i)_{Lc+s} + (l-L)c\lambda_i$ for all $l \geq L$ and for all i . Furthermore, $\lambda_{i_s} \geq \lambda_i$ for all i , and $\varepsilon \neq (h_{i_s})_{Lc+s} \geq (h_i)_{Lc+s}$ for all i with $\lambda_i = \gamma_s$. So if we define $K_s = L + \max\left(0, \max_{\substack{i \in N_s \\ \lambda_i \neq \gamma_s}} \left(\frac{(h_i)_{Lc+s} - (h_{i_s})_{Lc+s}}{c(\gamma_s - \lambda_i)}\right)\right)$ with $\max \emptyset = 0$ by definition, then we have $(h_{i_s})_{lc+s} \geq (h_i)_{lc+s}$ for all $l \geq K_s$ and all i . Hence, $g_{lc+s} = (h_{i_s})_{lc+s}$ for all $l \geq K_s$. As a consequence, (5) also holds for this case if we select $K \geq K_s$.

So if we define $K = \max(K_0, K_1, \dots, K_{c-1})$ then (5) holds for all s .

Assume $\bigoplus_{i=1}^m \lambda_i = \lambda_j$. Since $\lambda_j \neq \varepsilon$, there exists at least one index s^* such that $(h_j)_{Lc+s^*} \neq \varepsilon$.

Since $g_{Lc+s^*} = (h_{i_{s^*}})_{Lc+s^*}$ for all $l \geq K$ and since $\lambda = \lambda_{i_{s^*}}$ is the rate of $h_{i_{s^*}}$, i.e. the smallest λ for which (1) holds, $\gamma_s = \lambda_{i_{s^*}} = \lambda_j$ is also the smallest γ_s for which (5) holds. ■

Proof of Lemma 3.2: For sake of simplicity, we shall only prove the lemma for the case $m = 2$. The proof for $m > 2$ follows similar lines.

In this proof we always assume that $r, s \in \{0, \dots, c-1\}$, $p, q, i, k \in \mathbb{N}$.

Since h_1 and h_2 are ultimately geometric, there exists an integer L such that

$$(h_i)_{Lc+pc+s} = \lambda_i^{\otimes pc} \otimes (h_i)_{Lc+s} \quad \text{for all } p, r \text{ and } i = 1, 2 \quad (13)$$

(cf. (12) with $l = L + p$). We have

$$(h_1 \otimes h_2)_k = \bigoplus_{i=0}^{Lc+c-1} (h_1)_i \otimes (h_2)_{k-i} \oplus \bigoplus_{i=Lc+c}^{k-Lc-c} (h_1)_i \otimes (h_2)_{k-i} \oplus \bigoplus_{i=0}^{Lc+c-1} (h_1)_{k-i} \otimes (h_2)_i \quad (14)$$

for all $k \geq 2(Lc + c)$. Now we consider an arbitrary term of the second max-plus-algebraic sum of (14). Let $k \geq 2(Lc + c)$ and $i \in \{Lc + c, \dots, k - Lc - c\}$. Select p, q, r, s such that

$i = Lc + pc + r$ and $k - i = Lc + qc + s$. It is easy to verify that we have $\alpha^{\otimes p} \otimes \beta^{\otimes q} \leq \alpha^{\otimes p+q} \oplus \beta^{\otimes p+q}$ for all $\alpha, \beta \in \mathbb{R}_\varepsilon$ and all $p, q \in \mathbb{N}$. Hence,

$$\begin{aligned}
(h_1)_i \otimes (h_2)_{k-i} &= \lambda_1^{\otimes pc} \otimes (h_1)_{Lc+r} \otimes \lambda_2^{\otimes qc} \otimes (h_2)_{Lc+s} && \text{(by (13))} \\
&\leq \left(\lambda_1^{\otimes (p+q)c} \oplus \lambda_2^{\otimes (p+q)c} \right) \otimes (h_1)_{Lc+r} \otimes (h_2)_{Lc+s} \\
&\leq \lambda_1^{\otimes (p+q)c} \otimes (h_1)_{Lc+r} \otimes (h_2)_{Lc+s} \oplus \lambda_2^{\otimes (p+q)c} \otimes (h_1)_{Lc+r} \otimes (h_2)_{Lc+s} \\
&\leq (h_1)_{Lc+r} \otimes (h_2)_{Lc+(p+q)c+s} \oplus (h_1)_{Lc+(p+q)c+r} \otimes (h_2)_{Lc+s} && \text{(by (13)).}
\end{aligned}$$

Since $Lc+r \leq Lc+c-1$ and $Lc+r+Lc+s+(p+q)c = k$, the term $(h_1)_{Lc+r} \otimes (h_2)_{Lc+(p+q)c+s}$ also appears in the first max-plus-algebraic sum of (14). Similarly, it can be shown that $(h_1)_{Lc+(p+q)c+r} \otimes (h_2)_{Lc+s}$ also appears in the third max-plus-algebraic sum of (14). So the second max-plus-algebraic sum in (14) is redundant and can be omitted.

Now we define the sequences $f_{i,j}$ for $i = 0, \dots, Lc+c$ and $j = 1, 2$ with $(f_{1,i})_k = (h_1)_i \otimes (h_2)_{k-i}$ and $(f_{2,i})_k = (h_1)_{k-i} \otimes (h_2)_i$. The sequences $f_{i,j}$ are ultimately geometric with rate λ_j and cyclicity c_j . As shown above, the terms of $h_1 \otimes h_2$ coincide with $\bigoplus_{i,j} f_{i,j}$ for k large enough.

Therefore, we can now apply Lemma 3.1 provided that we select K such that $K \geq 2(Lc+c)$. ■

Note that in general we do not have $\gamma_s = \bigoplus_{i=1}^m \lambda_i$ for all indices s in Lemma 3.1 as is shown by the following example.

Example A.1 Consider the ultimately geometric sequences

$$\begin{aligned}
h_1 &= 0, \varepsilon, 1, 3, \varepsilon, 4, 6, \varepsilon, 7, 9, \varepsilon, 10, \dots \\
h_2 &= 0, \varepsilon, 0, \varepsilon, 0, \varepsilon, 0, \varepsilon, 0, \varepsilon, 0, \varepsilon, \dots
\end{aligned}$$

with $\lambda_1 = 1$, $c_1 = 3$, $\lambda_2 = 0$ and $c_2 = 2$. We have

$$h_1 \oplus h_2 = 0, \varepsilon, 1, 3, 0, 4, 6, \varepsilon, 7, 9, 0, 10, 12, \varepsilon, 13, 15, 0, 16, \dots$$

This sequence is ultimately periodic with $c = \text{lcm}(c_1, c_2) = \text{lcm}(2, 3) = 6$. Furthermore, the smallest γ_s s for which (5) holds are $\gamma_0 = \gamma_2 = \gamma_3 = \gamma_5 = 1$, $\gamma_1 = \varepsilon$ and $\gamma_4 = 0$. Note that $\varepsilon = \gamma_1 \neq \lambda_1 \oplus \lambda_2 = 1 \oplus 0 = 1$ and $\varepsilon = \gamma_1 \notin \{\lambda_1, \lambda_2\} = \{1, 0\}$. □